From CDC's 1994 School Health Policies and Programs Study (SHPPS)

About the School Health Policies and Programs Study (SHPPS)

SHPPS is a national survey periodically conducted to assess school health policies and programs at the state, district, school, and classroom levels. Results from the 1994 SHPPS were published in the <u>Journal of School Health</u>, Volume 65, Number 8, October 1995.

Alcohol and Other Drug (AOD)-Use Prevention Policy Characteristics

- 90% of states required districts or schools to have a policy that prohibits AOD use among students.
- 97% of districts and 97% of middle/junior and senior high schools had a written policy prohibiting AOD use among students.
- 64% of districts and 71% of middle/junior and senior high schools defined a drug-free school zone around school grounds.

Teacher and Staff Training on AOD Prevention

During the two years preceding the survey:

- 96% of states and 50% of school districts offered in-service training for teachers on AOD-use prevention.
- ◆ 23% of teachers* received training on AOD-use prevention.
- 72% of states and 39% of districts offered training on AOD-use prevention for school health services staff.
- ◆ 50% of school health services staff received training on AOD-use prevention.

AOD Rehabilitation Services

 48% of senior high schools and 39% of middle/junior high schools provided AOD rehabilitation services in or through the school.

Percentage of All States Recommending and All Districts Including Specific Statements, Rules, or Procedures in Policies Prohibiting Alcohol and Other Drug Use

| Statements, rules, or procedures | States | Districts |
|--|--------|-----------|
| Descriptions of violations and possible consequences | 82% | 85% |
| Procedures for communicating the policy to students, | | |
| staff, and parents/guardians | 82% | 74% |
| Support for prevention education | 82% | 72% |
| Procedures for implementing policy | 64% | 60% |

^{*}SHPPS interviewed both teachers of courses that focused primarily on health education topics, and teachers of courses that included some health education content but focused primarily on another subject (e.g., biology).

AOD-Use Prevention Education

Requirements


- ◆ 75% of states and 86% of districts required AOD-use prevention education.
- 90% of middle/junior and senior high schools included AOD-use prevention in a required course.

Topics Taught

- ♦ 70% of teachers taught AOD-use prevention as a major topic within their courses.
- ♦ 68% of teachers spent fewer than six class periods on AOD-use prevention.

Percentage of Middle/Junior and Senior High School Teachers Teaching Specific AOD-Use Prevention Topics

| Topic Percentage of teachers teac | |
|---|-----------|
| Short-term risks associated with alcohol use | 78% |
| Short-term risks associated with drug use | 75% |
| Effects of AOD use on decision making | 65% |
| Social influences on AOD use | 64% |
| Social norms towards risk behaviors related to AO | D use 60% |
| | |



For More Information

For additional information on SHPPS, contact the Centers for Disease Control and Prevention (CDC), National Center for Chronic Disease Prevention and Health Promotion, Division of Adolescent and School Health, 4770 Buford Highway, NE, Mailstop K-33, Atlanta, GA 30341-3717, telephone 770-488-3257, http://www.cdc.gov/nccdphp/dash